

**Table A.3B – Tobacco Product and Alcohol Use in the Past Month among Young Adults Aged 18 to 25**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
<b>TOBACCO PRODUCTS</b>	45.3* (0.48)	44.8* (0.48)	44.6* (0.50)	44.3* (0.48)	44.0* (0.49)	41.9* (0.50)	41.4* (0.47)	41.6* (0.50)	40.9* (0.49)	39.5* (0.49)	38.1* (0.47)	37.0* (0.49)	35.0* (0.54)	33.0* (0.48)	30.0 (0.48)
Cigarettes	40.8* (0.48)	40.2* (0.47)	39.5* (0.49)	39.0* (0.47)	38.5* (0.48)	36.2* (0.49)	35.7* (0.45)	35.8* (0.48)	34.3* (0.47)	33.5* (0.47)	31.8* (0.47)	30.6* (0.46)	28.4* (0.53)	26.7* (0.46)	23.5 (0.47)
Daily Cigarette Smoking <sup>1</sup>	51.8* (0.72)	52.7* (0.69)	51.6* (0.72)	50.1* (0.73)	48.8* (0.77)	49.2* (0.76)	47.8* (0.81)	45.3* (0.80)	45.8* (0.80)	45.3* (0.86)	45.1* (0.88)	43.1* (0.83)	43.0* (0.91)	42.0 (1.02)	39.9 (1.03)
Smoked 1+ Packs of Cigarettes per Day <sup>2</sup>	39.1* (0.93)	37.1* (0.88)	34.9* (0.86)	36.9* (0.93)	34.4* (0.93)	32.9* (0.92)	31.6* (0.91)	29.5 (0.92)	27.3 (0.94)	26.1 (0.97)	25.1 (0.90)	22.3* (0.90)	22.5* (1.16)	22.5* (1.11)	26.2 (1.42)
Smokeless Tobacco	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	5.4 (0.22)	5.2 (0.22)
Cigars	11.0* (0.27)	11.4* (0.26)	12.7* (0.30)	12.0* (0.28)	12.1* (0.29)	11.9* (0.28)	11.4* (0.29)	11.5* (0.29)	11.3* (0.30)	10.9* (0.29)	10.7* (0.27)	10.0* (0.29)	9.7* (0.30)	8.9 (0.27)	8.8 (0.27)
Pipe Tobacco	1.1* (0.08)	0.9* (0.08)	1.2* (0.09)	1.5 (0.11)	1.3* (0.10)	1.2* (0.10)	1.4 (0.10)	1.8 (0.12)	1.8 (0.12)	1.9 (0.14)	1.8 (0.11)	2.2* (0.14)	1.9 (0.13)	1.8 (0.13)	1.7 (0.12)
<b>ALCOHOL</b>	60.5* (0.53)	61.4* (0.50)	60.5* (0.51)	60.9* (0.51)	62.0* (0.51)	61.3* (0.52)	61.1* (0.49)	61.8* (0.52)	61.4* (0.50)	60.7* (0.54)	60.2* (0.49)	59.6* (0.53)	59.6* (0.56)	58.3 (0.53)	57.1 (0.55)
Binge Alcohol Use	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	39.0 (0.51)	38.4 (0.54)
Heavy Alcohol Use	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	10.9 (0.33)	10.1 (0.32)

nc = not comparable due to methodological changes.

NOTE: Estimates shown are percentages with standard errors included in parentheses.

\* The difference between this estimate and the 2016 estimate is statistically significant at the .05 level.

<sup>1</sup> Percentages for daily cigarette smoking are among past month cigarette smokers.

<sup>2</sup> Percentages for smoking one or more packs of cigarettes per day are among daily cigarette smokers in the past month. Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2016.

**Table A.4B – Tobacco Product and Alcohol Use in the Past Month among Adults Aged 26 or Older**

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
<b>TOBACCO PRODUCTS</b>	29.9* (0.44)	29.3* (0.41)	28.5* (0.41)	29.0* (0.43)	29.4* (0.43)	28.6* (0.42)	28.4* (0.44)	27.3* (0.40)	27.2* (0.42)	26.3* (0.41)	27.0* (0.42)	25.7* (0.40)	25.8* (0.33)	24.5 (0.32)	24.6 (0.33)
Cigarettes	25.2* (0.42)	24.7* (0.41)	24.1* (0.39)	24.3* (0.39)	24.7* (0.40)	24.1* (0.40)	23.8* (0.41)	23.0* (0.39)	22.8* (0.38)	21.9* (0.39)	22.4* (0.40)	21.6* (0.38)	21.5* (0.32)	20.0 (0.31)	20.2 (0.31)
Daily Cigarette Smoking <sup>1</sup>	68.8* (0.87)	68.0* (0.86)	67.8* (0.80)	68.9* (0.79)	67.9* (0.74)	66.3* (0.83)	67.0* (0.86)	67.2* (0.84)	64.8* (0.86)	66.5* (0.88)	66.0* (0.85)	64.9* (0.88)	63.3 (0.72)	62.7 (0.76)	62.2 (0.75)
Smoked 1+ Packs of Cigarettes per Day <sup>2</sup>	57.1* (1.12)	58.0* (0.99)	59.2* (1.05)	55.1* (1.02)	54.5* (1.00)	55.1* (1.06)	53.0* (1.10)	49.4* (1.16)	48.8* (1.09)	47.4* (1.05)	45.2 (1.09)	44.7 (1.15)	43.3 (0.93)	44.1 (0.98)	43.1 (1.00)
Smokeless Tobacco	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	3.2 (0.13)	3.1 (0.12)
Cigars	4.6 (0.18)	4.5 (0.18)	4.6* (0.17)	4.7* (0.18)	4.6* (0.18)	4.4 (0.16)	4.4 (0.18)	4.4 (0.18)	4.4 (0.17)	4.2 (0.18)	4.5 (0.19)	4.1 (0.17)	3.9 (0.12)	4.3 (0.14)	4.2 (0.13)
Pipe Tobacco	0.8 (0.09)	0.6 (0.07)	0.7 (0.08)	0.8 (0.08)	0.9 (0.09)	0.8 (0.09)	0.6 (0.07)	0.7 (0.07)	0.7 (0.07)	0.7 (0.07)	0.9 (0.09)	0.7 (0.07)	0.7 (0.06)	0.8 (0.06)	0.7 (0.06)
<b>ALCOHOL</b>	53.9 (0.53)	52.5* (0.49)	53.0* (0.51)	55.1 (0.51)	53.7 (0.49)	54.1 (0.52)	54.7 (0.50)	54.9 (0.48)	54.9 (0.48)	55.1 (0.49)	55.6 (0.48)	55.9* (0.50)	56.5* (0.39)	55.6 (0.38)	54.6 (0.38)
Binge Alcohol Use	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	24.8 (0.32)	24.2 (0.32)
Heavy Alcohol Use	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	6.4 (0.17)	6.0 (0.17)

nc = not comparable due to methodological changes.

NOTE: Estimates shown are percentages with standard errors included in parentheses.


\* The difference between this estimate and the 2016 estimate is statistically significant at the .05 level.

<sup>1</sup> Percentages for daily cigarette smoking are among past month cigarette smokers.

<sup>2</sup> Percentages for smoking one or more packs of cigarettes per day are among daily cigarette smokers in the past month. Respondents with missing data for number of cigarettes smoked per day were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2016.

**Figure 14. Current Alcohol Use among People Aged 12 to 20: Percentages, 2002-2016**



<sup>+</sup> Difference between this estimate and the 2016 estimate is statistically significant at the .05 level.

**Figure 14 Table. Current Alcohol Use among People Aged 12 to 20: Percentages, 2002-2016**

Use	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16
Current	28.8 <sup>+</sup>	29.0 <sup>+</sup>	28.7 <sup>+</sup>	28.2 <sup>+</sup>	28.4 <sup>+</sup>	28.0 <sup>+</sup>	26.5 <sup>+</sup>	27.2 <sup>+</sup>	26.2 <sup>+</sup>	25.1 <sup>+</sup>	24.3 <sup>+</sup>	22.7 <sup>+</sup>	22.8 <sup>+</sup>	20.3	19.3

<sup>+</sup> Difference between this estimate and the 2016 estimate is statistically significant at the .05 level.